

The eclipsing Intermediate Polar V597 Pup (Nova Puppis 2007)

Brian Warner^{*}, Patrick A. Woudt[†]

Department of Astronomy, University of Cape Town, Private Bag X3, Rondebosch 7701, South Africa

2009 April 11

ABSTRACT

Photometric observations of V597 Pup made in 2008, 9.1 mag below maximum, 4 months after eruption, showed no certain orbital modulation but exhibited a quintuplet of oscillations centred on a period 261.9 s and uniform splitting at a frequency $\sim 2.68 \text{ h}^{-1}$. One year later the system had fallen in brightness by a further 2.5 mag, showed deep eclipses with a period of 2.6687 h, and the 261.9 s modulation at a reduced amplitude. There is often power near the ‘subharmonic’ at 524 s, showing that the shorter periods observed are actually first harmonics.

V597 Pup is thus an intermediate polar and is in the ‘orbital period gap’. Furthermore it is the first to show a prominent secondary eclipse, caused by passage of the optically thick disc in front of the irradiated side of the secondary star.

Key words: binaries – close – novae – stars: oscillations – stars: individual: V597 Pup, cataclysmic variables

1 INTRODUCTION

V597 Pup was discovered as a nova at $V \sim 7.5$ on 14 November 2007 (Pereira, McGaha & Rhoades 2007), reaching $V \sim 6.2$ at maximum and declining smoothly for at least 5 magnitudes as a very fast nova, with $t_2 = 2.5 \text{ d}$ (Naik, Banerjee & Nashok 2009), which is nearly the shortest recorded (only V838 Her and MU Ser are shorter: Table 5.2 of Warner 1995). Its infrared red spectra and X-Ray emission (as a super-soft source) in early stages of decline are discussed by Naik et al., who classify it as a He/N nova and deduce that it has a white dwarf mass close to the Chandrasekhar limit.. By March 2008 it had reached $V \sim 15.3$; our later observations place it (out of eclipse) at ~ 17.7 in February 2009 and ~ 17.9 a month later. No post-eruption spectra have been published. A possible pre-eruption image at $V \sim 20$ is present on the Digitized Sky Survey (Pereira et al. 2007). A finding chart prepared from one of our CCD images is given in Fig. 1. The position of V597 Pup appears to coincide with the proposed pre-eruption image.

In Section 2 we give the results and an initial analysis of our observations. In Section 3 we look at the overall geometry of V597 Pup. Section 4 expands on the analysis by treating V597 Pup as an eclipsing intermediate polar, and Section 5 summarizes our interpretation of this interesting cataclysmic variable.




Figure 1. CCD image of V597 Pup (indicated by the markers) taken on 2009 March 18 (out of eclipse). The field of view is 50 by 34 arcsec, north is up and east is to the left.

2 OBSERVATIONS


We used the University of Cape Town (UCT) CCD photometer, as described by O’Donoghue (1995), in frame transfer mode and with white light, on the 1.9-m (74-in) reflector at the Sutherland site of the South African Astronomical Observatory (SAAO). Tab. 1 contains a detailed observing log. Our magnitude scale was derived using hot white dwarf standards, but because of the non-standard spectral distributions of the spectrum of the nova remnant, and the use

^{*} email: Brian.Warner@uct.ac.za

[†] email: Patrick.Woudt@uct.ac.za

Table 1. Observing log of photometric observations

Run	Date of obs. (start of night)	HJD of first obs. (+2450000.0)	Length (h)	t_{in} (s)	V (mag)
S7807	2008 Mar 15	4541.26125	2.77	6	15.2
S7808	2008 Mar 16	4542.24472	3.53	6	15.2
S7810	2008 Mar 17	4543.27429	3.82	6	15.3
S7812	2008 Mar 18	4544.25449	3.30	6	15.3
S7828	2009 Feb 25	4888.26865	7.30	30	17.8 [†]
S7830	2009 Feb 26	4889.28184	6.95	30	17.7 [†]
S7832	2009 Feb 27	4890.26684	7.39	30	17.7 [†]
S7834	2009 Feb 28	4891.25647	4.29	30	17.7 [†]
S7840	2009 Mar 03	4894.30579	1.15	30	17.7 [†]
S7842	2009 Mar 18	4909.24058	5.42	30	17.9 [†]
S7843	2009 Mar 19	4910.23990	5.51	30	17.9 [†]
S7856	2009 Mar 23	4914.29437	1.68	30	17.9 [†]



[†] Out of eclipse.**Figure 2.** The 2008 light curves of V597 Pup phased on the orbital ephemeris given in Eq. 1. The light curves of runs S7808, S7810, S7812 have been displaced by 0.4, 0.8 and 1.2 mag, respectively, for display purposes.

of white light, our magnitudes approximate a V scale only to ~ 0.1 mag.

2.1 Light curves on the orbital time scale

The light curves from the 2008 data are shown in Fig. 2, aligned in phase with the orbital period that we deduce below. A repetitive hump with an amplitude ~ 0.2 mag is seen, but no obvious eclipse features; unfortunately our light curves had lengths not much greater than the (then unknown) orbital period. The Fourier transform (FT) of the combined light curves is shown in Fig. 3. The highest peak has a frequency of $91.8 \mu\text{Hz}$, but there are aliases of almost equal amplitude either side at 80.3 and $103.3 \mu\text{Hz}$ (periods of 3.46 , 3.03 and 2.69 h). There is also a set of frequencies which could be first harmonics, at 173.7 , 185.2 , 196.7 and $208.1 \mu\text{Hz}$. The uncertainties on these figures are $\sim 0.5 \mu\text{Hz}$.

Light curves from the 2009 observations are shown in Fig. 4, where it is clear that the drop in system brightness by ~ 2.5 mag over one year has revealed broad eclipses ~ 0.6 mag deep with respect to the out-of-eclipse level away

**Figure 3.** The low frequency Fourier transform of the combined 2008 observations of V597 Pup.**Figure 4.** The 2009 light curves of V597 Pup phased on the orbital ephemeris given in Eq. 1. The light curve of run S7828 is displayed at the correct magnitude, the others have been displaced vertically for display purposes only.

from the reflection region (see below). The fundamental and harmonics in the FT of these light curves provide an orbital period of 2.6687 h ($104.089 \mu\text{Hz}$), which shows that the orbital modulation, with first harmonic, was present in the 2008 light curves. The ephemeris for minimum light is

$$\text{HJD}_{\min} = 2454888.3280 + 0^{\text{d}}11119 \text{ E.} \quad (1)$$

If the drop in brightness (by a factor ~ 10) between the two sets of observations was due largely or entirely to the dispersal and cooling of nova ejecta far from the central binary, the eclipses and their related humps could well have been present even in the earlier light curves – there are some features that resemble such diluted structures.

The 2009 light curves for February and March averaged over the binary period are seen in Fig. 5 and show repetitive structures between the primary eclipses. Remembering that these are of a fast nova only 15 months after outburst we would perhaps expect the white dwarf primary still to be very hot, so that at the small distance from the secondary implied by the short orbital period a strong reflection effect




Figure 5. The average light curves of V597 Pup for the February and March 2009 observations, plotted separately.




Figure 6. The average light curves of V597 Pup as in Fig. 5 - the 2009 March light curve is shifted vertically by 0.18 mag to match the 2009 February average magnitude - with overplotted the expected sinusoidal variation of a strong reflection effect (dashed line) of ~ 0.54 mag peak-to-peak amplitude. The depth of the deepest secondary eclipse (S7842 - see Fig. 4) is an indication of the approximate minimum brightness at secondary eclipse, marked by the horizontal dotted line.

could be present. Between eclipses there is indeed what may be described as a reflection effect which would be 0.3 or 0.4 mag in amplitude were it not suppressed either side of phase 0.5 by a secondary eclipse of total width equal to that of primary eclipse. In Fig. 6 we sketch what we suggest would be the reflection effect without any secondary eclipse; the asymmetry, namely the early fall in brightness before the start of primary eclipse is probably caused by partial obscuration of the bright face of the companion star by the gas stream passing from it to the outer edge of the disc. The reflection effect and secondary eclipse are similar, but of greater amplitude, to what we saw in the nova remnant DD Cir (Woudt & Warner 2003), where we ascribed a shallow secondary eclipse to an optically thick almost edge-on disc passing across the heated hemisphere of the companion star. In V597 Pup the total widths of the primary and secondary eclipses are both $\sim \pm 0.15$ of P_{orb} .

There is great variability in the profile and depth of the secondary eclipse, as could be expected from a disc of rapidly changing structure. Figs. 5 and 6 show that the 0.2 mag decrease in brightness from February to March 2009 has resulted in a deeper primary eclipse as the fading ejecta reduce the amount of in-fill.

Inspection of the individual light curves in Fig. 4 shows that during primary and secondary eclipses there is flickering similar to that seen out of eclipse, so at least one flick-




Figure 7. The high frequency Fourier transform of the combined 2008 observations of V597 Pup (top panel). The middle panel shows the detailed structure of the quintuplet around $3818 \mu\text{Hz}$, where the vertical bars correspond to the frequencies listed in Tab. 2. The bottom panel shows the FT around $3818 \mu\text{Hz}$ after pre-whitening (with a simultaneous 5 sinusoid least squares fit) with the frequencies listed in Tab. 2.

ering source is still visible during eclipse. Apart from that, certainly the secondary and possibly the primary eclipse, appear flat bottomed, as in a total eclipse or a transit.

2.2 Short period optical oscillations

The FT of the combined 2008 light curves is shown in Fig. 7. There is red noise at low frequencies, associated with flickering and with harmonics of the orbital modulation, but an isolated cluster of peaks in the vicinity of $3800 \mu\text{Hz}$ is evident. On detailed examination five components are found, with the frequencies and amplitudes listed in Tab. 2, found from a 5 sinusoid simultaneous least squares fit. Uncertainties of the frequencies are all $\sim \pm 0.5 \mu\text{Hz}$ and of the amplitudes are ± 0.7 mmag. The phases (with arbitrary zero point) are quoted as fractions of cycles. The largest amplitude signal is at 261.9 s.

As further evidence for the significance of these signals we produce in Fig. 8 an O - C phase diagram relative to the central frequency and comparison with the same diagram produced by a simulation using the five sinusoidal signals listed in Tab. 2.

The mean splitting between the components in 2008 is $104.15 \mu\text{Hz}$, which, within errors, is the same as the orbital frequency. It is therefore clear that in 2008 a quintuplet of frequencies was present with internal splitting equal to the orbital frequency Ω , which is the characteristic signature of an intermediate polar (IP) where the white dwarf primary's frequency ω , and/or its reprocessed signal $\omega - \Omega$, acquires orbital sidebands. This occurs through reprocessing of a ro-




Figure 8. Phase variations (observed – calculated) of the 3818 μHz signal in the 2008 data (observed: top panel; simulated (see text for details): bottom panel). Different symbols represent data from different nights. Each dot represents ~ 6 cycles of the 3818 μHz modulation, with a 50% overlap.

Table 2. The high frequencies (first harmonics) in V597 Pup.

2008			2009 February	
Frequency (μHz)	Ampl. (mmag)	Phase	Frequency (μHz)	Ampl. (mmag)
3610.0	2.2	-0.05 ± 0.09		
3713.4	2.0	-0.37 ± 0.10	3714.6	2.7
3818.2	3.4	-0.21 ± 0.06	3817.8	6.3
3922.4	1.6	-0.45 ± 0.12		
4026.8	2.3	$+0.24 \pm 0.08$	4027.3	3.1

tating beam of high energy radiation, emitted from the accretion zone(s) on the primary, off regions of varying cross section or varying visibility rotating in orbit (Warner 1986). This shows that the central binary was free from obscuration during the 2008 observations and therefore that much of the luminosity came from distant ejecta and not an optically thick wind close to the primary. The apparent amplitude of the oscillations should be increased by a factor of at least 10 to remove the diluting effect of these bright ejecta, so relative to the central binary luminosity the amplitudes are ~ 0.03 mag. There is no significant extra power at the harmonics or subharmonic of the 261.9 s signal.

In the February 2009 light curves the sidebands are largely conflated with noise in the FT, but the central modulation is very clear – see Fig. 9 – and near numerical coincidences with the 3713.4 and 4026.8 μHz sidebands are present. The amplitude, relative to the brightness of the central binary, has dropped by a factor ~ 5 . The average of the two years’ frequency determinations is 3818.0 μHz , or a period of 261.9 s. In the March 2009 light curves no components of the 2008 quintuplet are definitely present, but both February and March 2009 light curves possess signals near ‘subharmonics’ of the quintuplet: see Fig. 10. These are listed in Tab. 3. In February 2009 the strongest signal was at 1907.5 μHz (2×262.1 s), with a nearby 1-d alias; in March 2009 there was a large amplitude window pattern which included the 1907.5 μHz modulation as one of the aliases. The clear presence of a subharmonic to the 261.9 s signal shows that the periods listed in Tab. 2 are almost certainly first harmonics, not fundamentals.




Figure 9. The high frequency Fourier transform of the combined 2009 February observations of V597 Pup around 3818 μHz . The vertical bars correspond to the frequencies listed in Tab. 2.




Figure 10. The high frequency Fourier transform of V597 Pup around 1905 μHz of the combined 2009 February observations (top panel), and the combined 2009 March observations (lower panel). The vertical bars correspond to the selected frequencies listed in Tab. 3.

Table 3. The high frequencies (fundamentals) in V597 Pup.

2009 February		2009 March	
Frequency [†] (μHz)	Ampl. (mmag)	Frequency [†] (μHz)	Ampl. (mmag)
		<i>1930.8 $\pm$ 0.3</i>	<i>11.3 $\pm$ 2.8</i>
		<i>1919.2 $\pm$ 0.3</i>	<i>11.1 $\pm$ 2.8</i>
		<i>1909.9 $\pm$ 0.3</i>	<i>9.2 $\pm$ 2.8</i>
1907.5 $\pm$ 0.5	8.1 $\pm$ 0.5	1907.5 $\pm$ 0.6	8.9 $\pm$ 2.0
1896.1 $\pm$ 0.5	8.0 $\pm$ 0.5		

[†] Aliases are given in italics.

As is usual with IPs (Warner 2004) we found no evidence for dwarf nova oscillations in the FTs of V397 Pup.

3 THE STRUCTURE OF V597 PUP

The drop in luminosity from 2008 to 2009 revealed an IP in the orbital period gap (Warner 1995), with a rotation period of 8.7 min. The reduction in amplitudes of the spin modulations over the year is probably the result of diminishing $\dot{M}$ as the primary cools and lessens its irradiation of the secondary.

It is not possible, with the flickering and variations from orbit to orbit, to give precise values for eclipse ingress and egress timings, but we can show the effect in the light curve of the various components. First, a simple calculation shows that the measured widths of the eclipses imply an unusually large disc radius. From the expected high mass for a very fast nova, and the mass and radii relationships for CV secondaries (Warner 1995), we adopt $M(1) = 1.3 M_{\odot}$, $M(2) = 0.22 M_{\odot}$, $R(2) = 1.9 \times 10^{10}$ cm. These lead to a separation $a = 7.82 \times 10^{10}$ cm, a radius $R(L)$ of the Roche lobe of the primary of 4.2×10^{10} cm, and expected radius of a high $\dot{M}$ disc of $R_d = 0.7R(L) = 2.9 \times 10^{10}$ cm. The first and last contact phases $\pm\phi$ of eclipses, relative to phase 0, assuming an inclination near 90° , is found from $\sin\phi = \frac{[R(2)+R]}{a}$, where R is selected from above. For the usual $R = R_d$ we find $\phi = 0.10$, but for $R = R(L)$ we have $\phi = 0.14$, which is close to the observed $\phi \sim 0.15$. The disc in V597 Pup fills the primary's Roche lobe to its maximal possible extent. The radius of the disc is considerably larger than that of the secondary star, which results in the flat bottomed secondary eclipse.

The phases of first and last contacts in the secondary eclipse, as the secondary star (made visible by irradiation from the hot primary) is eclipsed by the disc, relative to phase 0.5, are also $\pm\phi$, and the phases between second and third contact are $\pm(\phi - \theta)$, where $\sin\theta = \frac{[R(L)-R(2)]}{a}$, which are ± 0.09 , in reasonable agreement with what is observed for the emergence of the secondary, but immergence takes longer than predicted, possibly because at that phase the illuminated gas stream is also being eclipsed by the disc.

The general agreement between observed and expected time scales of the secondary eclipses adds weight to our interpretation of this unusual light curve.

With the short orbital period, and mass ratio $M_2/M_1 = 0.22$ adopted here, we might have expected an elliptical disc and superhumps to be present (see Chapter 3 of Warner 1995), but there is no evidence for them in the light curves and FTs. The reason may be that with an accretion disc of such large radius, greatly exceeding the 3:1 resonance radius that excites ellipticity, the resonance is ineffectual: Osaki & Meyer (2003) have shown how in a disc of large radius the 2:1 Lindblad resonance can suppress the 3:1 resonance. Our estimate of $\phi \sim 0.15$ in fact puts the outer edge of the accretion disc very close to the 2:1 resonance radius.

4 V597 PUP AS AN INTERMEDIATE POLAR

Mukai (2008) lists over 30 bona fide IPs, among which there are none with relatively rapid rotations that possess deep

eclipses suitable to provide detailed information about the structure of an IP. Eclipses are seen in XY Ari but it is an X-Ray source, hidden behind a dust cloud; DQ Her is far more favourable for observation but is not ideal as it has an inclination so high that much of the central region of the disc is hidden from view; DD Cen (Woudt & Warner 2003) is too faint to study fully. Thus V597 Pup is the first relatively standard IP found to have eclipses – which can be expected to grow a little deeper over the next few years as the residual emission from the ejecta decays. We analyse our observations further in order to develop a model for the system.

4.1 Interpretation of the Quintuplet in the 2008 Light Curves

The standard interpretation of multiple sidebands in IPs (Warner 1986) ascribes the (often very strong) $\omega - \Omega$ component to reprocessing of ω from the secondary star or the disc thickening near the bright spot or the inner Lagrangian point (where in such a large disc the vertical gravity is very low)¹, all of which may vary in cross section at frequencies Ω and 2Ω , and can thus generate $\omega - 2\Omega$, ω and $\omega + \Omega$ signals, as well as $\omega - \Omega$. This gives only four components, a quadruplet, whereas in V597 Pup we uniquely see a quintuplet. We suggest that the clue to understanding this comes from the unusual presence of a secondary eclipse: the $\omega - \Omega$ component is therefore modulated not only at Ω (with maximum at superior conjunction and minimum at inferior conjunction of the secondary), but also at 2Ω (another minimum being imposed by the secondary eclipse, where otherwise a maximum of reprocessing would occur). This generates $\omega - \Omega \pm 2\Omega$ components, so together with the standard quadruplet, the complete set will be $\omega - 3\Omega$, $\omega - 2\Omega$, $\omega - \Omega$, ω , $\omega + \Omega$. From what we deduced in Section 2.2 the 2008 quintuplets must be the first harmonics of then unobservable fundamentals.

This interpretation attributes the highest amplitude component, at 3818.0 μHz , to $2(\omega - \Omega)$, not to the harmonic of the primary's spin, 2ω . Interestingly, the well known 71 s modulation in the nova remnant DQ Her, which is also a high inclination binary, has recently been re-interpreted (Saito & Baptista 2009) as $(\omega - \Omega)$ and not ω , being a result of reprocessing from the thickened disc region around the bright spot.

4.2 The Phase Variation of the 262 s modulation through Primary Eclipse

The first phase shift discovered in the eclipse of an IP was for DQ Her (Warner et al. 1972), studied in great detail by Patterson, Robinson & Nather (1978). The phase (observed – calculated) of the 71 s modulation increases by $\sim 90^\circ$ from the beginning to mid-point of eclipse, jumps through $\sim 180^\circ$ and recovers from mid- to end of eclipse. By contrast, the stable Dwarf Nova Oscillations (DNOs) in the nova-like variable UX UMa decrease through 360° during eclipse (Nather & Robinson 1974); the same behaviour has been seen in HT

¹ The reader is reminded that sinusoidal modulation of ω by Ω generates signals at ω and $\omega \pm \Omega$.




Figure 11. The amplitude (upper panel) and phase variation (observed – calculated, lower panel) of the 3818 μHz modulation as a function of orbital phase (Eq. 1).

Cas (for a review of eclipse phase changes see Chapter 8 of Warner 1995).

The phase changes have been successfully modeled by the reprocessing of high energy radiation emitted from accretion zones on the rotating primary (Chester 1979; Petterson 1980), and are a function of the relative amounts of direct radiation from the primary and those from the front and back portions of the accretion disc.

In V597 Pup the phase behaviour of the 262 s signal in 2009 is of a kind that has not been observed before. Our initial impression was of a 360° increase in phase during eclipse – which might be thought to indicate that the primary rotates retrogradely in the system. But further investigation showed that Chester (1979) had anticipated the situation – he pointed out that if the phase of the first harmonic is measured in a high inclination system (where only the beam sweeping across the back surface of the disc is observable, as in DQ Her) then a 180° increase would be seen, followed by an instantaneous decrease of 360° and a recovery increase of 180° . There is of course an ambiguity in how to plot such observed phase shifts, but we have chosen to use Chester’s insight rather than claim a retrogradely rotating primary. The phase variations through eclipse are present in all the high quality light curves; these and their average are shown in Fig. 11.

There is no significant reduction in amplitude near phase 0, which may be the result of the $\omega - \Omega$ signal having at least three sources (secondary and two regions of disc thickening). Furthermore, unlike DQ Her, where the radii of the secondary and disc are almost equal (Petterson 1980), in V597 Pup the secondary is considerably smaller and leaves much of the disc uncovered at phase 0.

As seen in Fig. 11, there are no significant phase changes associated with the secondary eclipse.




Figure 12. The amplitude and phase variation (upper two panels, respectively) of the 524 s modulation in run S7842. The lower two panels show the amplitude and phase variation of the 524 s modulation in run S7843. Each dot represents ~ 2 cycles of the 524 s modulation, with a 50% overlap. In the phase diagrams, only those points are plotted where the amplitude of the modulation is larger than 10 mmag.

4.3 The Phase Variation of the 524 s modulation

The phase and amplitude variations of the 1907.5 μHz (524 s) fundamental modulation seen in our best light curves (runs S7842 and S7843 of March 2009, both of which cover two orbits) are shown in Fig. 12. The results through the eclipses are not clear cut, which may result from varying contributions of different sources to the $\omega - \Omega$ signal, but is largely due to the modulation period being comparable to the ingress and egress durations.

However, there is a DQ Her-like variation of phase centred on the first primary eclipse of S7842, not repeated in the second eclipse, and a suggestive rise in the second eclipse of S7843. In addition there is a reverse phase change (also $\sim 180^\circ$) in the secondary eclipses of S7842 and a probably similar effect in the second secondary eclipse of S7843 (in the first of the secondary eclipses of this run the amplitude of the signal is quite low). There are also some indications of lower amplitudes at the times of both primary and secondary eclipses.

5 CONCLUSION

V597 Pup is unique in several ways and will repay further photometric and spectroscopic investigation before it fades

another two magnitudes to its pre-eruption brightness. It is the first cataclysmic variable found to have a deep secondary eclipse – the result of a background generated by the temporary extra brightness of one face of the otherwise low luminosity secondary star. Detailed study of this eclipse could reveal more about the structure of the outer parts of the accretion disc.

As an intermediate polar V597 Pup offers a rare opportunity to study reprocessing in a very high inclination system. It has some parallels with DQ Her, but the fundamental and first harmonics of the rotation and reprocessed signals are more complex.

ACKNOWLEDGMENTS

Our research is supported by the University of Cape Town and by the National Research Foundation.

REFERENCES

- Chester T.J., 1979, ApJ, 230, 167
Mukai K., 2008,
<http://asd.gsfc.nasa.gov/Koji.Mukai/iphome/iphome.html>
Naik S., Banerjee D.P.K., Ashok N.M., 2009, MNRAS, 394, 1551
Nather R.E., Robinson E.L., 1974, ApJ, 199, 637
O'Donoghue D., 1995, Baltic Ast, 4, 517
Osaki Y., Meyer F., 2003, A&A, 401, 325
Patterson J., Robinson E.L., Nather R.E., 1978, ApJ, 224, 570
Pereira A.J.S., McGaha J.E., Rhoades H., 2007, IAUC, 8895
Petterson J.A., 1980, ApJ, 241, 247
Saito R.K., Baptista R., 2009, ApJ, 693, L16
Warner B., 1986, MNRAS, 219, 347
Warner B., 1995, Cataclysmic Variable Stars, Cambridge Univ. Press, Cambridge
Warner B., 2004, PASP, 116, 115
Warner B., Peters W.L., Hubbard W.B., Nather R.E., 1972, MNRAS, 159, 321
Woudt P.A., Warner B., 2003, MNRAS, 340, 1011